

The Canadian Field-Naturalist

SUPPLEMENTARY MATERIAL:

Comparative reproductive parameters of sympatric Lesser Scaup (*Aythya affinis*) and Ring-necked Duck (*Aythya collaris*) in parkland Manitoba

GORDON S. HAMMELL, HOWARD V. SINGER, and LLWELLYN M. ARMSTRONG

Table S1. Assigned scores for local spring wetland condition and selected Lesser Scaup (*Aythya affinis*) reproductive periods (with corresponding change in pond water level), 2009–2018, Erickson, Manitoba.

Year	Spring wetland score	Pre-nest period		Nesting period		Total wetland score
		cm	Score	cm	Score	
2009	3.0	-3.3±0.8*	2	-16.5±1.1	0	5.0
2010	3.0	14.2±2.7	5	0.7±3.0	3	11.0
2011	5.0	3.5±1.9	3	-25.8±1.9	0	8.0
2012	0.5	2.9±0.7	3	-11.3±0.9	0	3.5
2013	3.0	-1.8±0.7	2	2.1±1.8	3	8.0
2014	4.0	0.5±1.9	3	-0.1±3.3	2	9.0
2015	2.5	-5.1±0.6	1	-2.8±1.0	2	5.5
2016	2.5	-8.8±1.7	1	-0.3±1.4	2	5.5
2017	4.0	-5.7±0.5	1	-11.8±0.7	0	5.0
2018	2.2	1.5±0.7	3	-9.8±1.6	1	6.2

*A negative value (±SE) represents a drop in water level. Levels were measured at 15 class IV and V wetlands on or near the block study area